

- KROODSMA, D.E. 1981. Ontogeny of bird song. Pages 518–532 in K. Immelmann, G. Barlow, L. Petrinovich, and M. Main [EDS.], *Early development in man and animals. The Bielefeldt project*. Cambridge Univ. Press, Cambridge, U.K.
- . 1984. Songs of the Alder Flycatcher (*Empidonax alnorum*) and Willow Flycatcher (*Empidonax traillii*) are innate. *Auk* 101:13–24.
- MIKKOLA, H. 1983. Owls of Europe. T. & A.D. Poyser, Calton, U.K.
- MYSTERUD, I. AND H. DUNKER. 1982. Food and nesting ecology of the Eagle Owl, *Bubo bubo* L., in four neighbouring territories in southern Norway. *Swedish Wildlife Research, Viltrevy* 12:3.
- PENTERIANI, V. 1996. The Eagle Owl. Calderini Ed., Bologna, Italy.
- AND F. PINCHERA. 1991. Comparison of the playback and the listening methods in the census of Eagle Owl (in Italian). Pages 385–388 in M. Fasola [ED.], *Atti II Seminario Italiano Censimenti Faunistici dei Vertebrati*. Suppl. Ric. Biol. Selv. Vol. XVI, Bologna, Italy.
- RALPH, C.J. AND J.M. SCOTT. [EDS.]. 1981. Estimating the number of terrestrial birds. *Stud. Avian Biol.* No. 6.
- , J.R. SAUER AND S. DROEGE. [EDS.]. 1995. Monitoring bird populations by point counts. Gen. Tech. Rep. PSW-GTR-149. Albany, CA, U.S.A.
- REID, J.R., R.B. HORN, AND E.D. FORSMAN. 1999. Detection rates of Spotted Owls based on acoustic-lure and live-lure surveys. *Wildl. Soc. Bull.* 27:986–990.
- SMITH, D.G. 1987. Owl census techniques. Pages 304–309 in R.W. Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre [EDS.], *Biology and conservation of northern forest owls: symposium proceedings*. Gen. Tech. Rep. RM-142, Fort Collins, CO U.S.A.
- SOKAL, R.R. AND F.J. ROHLF. 1995. Biometry. The principles and practice of statistics in biological research, 3rd ed. W.H. Freeman and Company, New York, NY U.S.A.
- STEWART, A.C., R.W. CAMPBELL, AND S. DICKIN. 1996. Use of dawn vocalizations for detecting breeding Cooper's Hawks in an urban environment. *Wildl. Soc. Bull.* 24:291–293.
- VOOUS, K.H. 1988. Owls of the Northern Hemisphere. William Collins Sons & Co., London, U.K.

Received 2 December 1999; accepted 20 March 2000

J. Raptor Res. 34(3):235–237

© 2000 The Raptor Research Foundation, Inc.

FOOD HABITS OF THE STRIPED OWL (*ASIO CLAMATOR*) IN BUENOS AIRES PROVINCE, ARGENTINA

JUAN P. ISACCH,¹ MARÍA S. BÓ, AND MARIANO M. MARTÍNEZ²

Laboratorio de Vertebrados, Departamento Biología, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata, Funes 3250, (7600) Mar del Plata, Argentina

KEY WORDS: *Striped Owl*; *Asio clamator*; *food habits*; *Argentina*.

The Striped Owl (*Asio clamator*) is a widespread species from Mexico through tropical and subtropical South America to Argentina (Grossman and Hamlet 1964, Canevari et al. 1991, Holt et al. 1999). It inhabits deciduous seasonal forests, lowland seasonal forests, gallery forests, lowland savannas, desert forests, and grasslands (Grossman and Hamlet 1964, Canevari et al. 1991, Holt et al. 1999). Despite its widespread distribution, the status of the Striped Owl is poorly known (Burton 1973, Holt et al. 1999) and it is con-

sidered to be a rare species in Buenos Aires Province in Argentina (Narosky and Di Giacomo 1993).

Studies of the Striped Owl in Argentina have focused mainly on anecdotal aspects of its biology and breeding ecology (e.g., Bledinger et al. 1987, Martínez et al. 1996). Its diet is poorly studied but the limited information that is available indicates that it preys mainly on small mammals (Grossman and Hamlet 1964, Burton 1973, Phelps and Meyer de Schauensee 1994) followed by birds, reptiles, and insects (Holt et al. 1999). Here, we report on the diet of Striped Owls in the southernmost extreme of its distribution in the southeastern portion of Buenos Aires Province, Argentina.

METHODS

Our study was carried out in Mar Chiquito Biosphere Reserve (37°40'S, 57°23'W), Buenos Aires Province, Argentina. The reserve covers 30 000 ha and supports a diverse array of habitats including ponds, salt marshes,

¹ Present address: Comisión de Investigaciones Científicas de la Provincia de Buenos Aires, Buenos Aires, Argentina.

² Deceased.

grasslands, woodlands, exotic tree plantations, and agricultural fields.

We found a pair of Striped Owls in an area of tala (*Celtis tala*) forest at Nahuel Rucá Ranch. The tala forest is derived from thorn forests (Espinal) and this relict patch represents the southernmost extreme of Espinal forest in Argentina. The patch covered a 6 ha area and was surrounded by a pond and grazed native grassland. There were a few houses and a plantation of eucalyptus (*Eucalyptus* spp.) trees nearby.

From August to November 1997, pellets and prey remains were collected in different plucking stations and roosting sites of the owls. Bird, mammal, and insect remains were identified based on bones, feathers, bills, hair, dentaries, and exoskeletons, and compared with the collections of Laboratorio de Vertebrados, Facultad de Ciencias Exactas y Naturales-Universidad Nacional de Mar del Plata and Museo Municipal de Ciencias Naturales “Lorenzo Scaglia” de Mar del Plata. The majority of prey were identified to species level. Bird and mammal weights were obtained from the literature (Salvador 1988, Camperi 1992, Redford and Eisenberg 1992) and unpublished data (M. Kittlein pers. comm.). A weight of 1 g was assigned to each insect prey species (Jiménez 1993).

RESULTS AND DISCUSSION

A total of 56 prey items was identified from 34 pellets and 3 prey remains (pile of feathers). Rodents were the main prey (55.4%) followed by birds (42.9%) and insects (1.8%, Table 1). *Rattus* spp. was the most common prey item (43%). Among birds, Rufous-collared Sparrows (*Zonotrichia capensis*, 23.2%) and Eared Doves (*Zenaida auriculata*, 17.9%) were most frequently taken. Other items comprised only a small fraction of the diet (16.2%).

Prey weights ranged from a low of 1 g in the case of insects to a high 630 g in the case of *Cavia aperea* (Table 1). Rodents comprised up to 81.5% of prey by weight and *Rattus* spp. contributed with the highest value (66.9%) followed by *Cavia aperea* (14%). The occurrence of adult *C. aperea* in the diet was surprising, since they weigh more than one and a half times as much as Striped Owls (maximal weight recorded of Striped Owl is 485 gr, Salvador 1988). We were not certain if *C. aperea* were eaten as carrion or actually hunted but Striped Owls are highly adapted to hunt live prey (Holt et al. 1999). The biomass contribution of birds was minor (20.1%) with Eared Doves contributing the largest amount (12%, Table 1). Other studies have confirmed that birds are common in the diets of Striped Owls (Grossman and Hamlet 1964, Burton 1973, Phelps and Meyer de Schauensee 1994, Holt et al. 1999).

Our results agree with those of Martínez et al. (1996) who studied the diet of Striped Owls in an area of shrub and exotic trees in Laguna de Los Padres Reserve, located 35 km north of Nahuel Ruca in Buenos Aires Province. They recorded seven bird and mammal species in the diet; three of which (*Reithrodon auritus*, *Holochilus brasiliensis*, and *Carduelis magellanica*) were absent in the diet of the Striped Owls we studied.

Table 1. Frequency of prey items, weight of individual prey and total percent biomass of prey in the diet of Striped Owl (*Asio clamator*) in Mar Chiquito Biosphera Reserve, Buenos Aires Province, Argentina.

PREY	FRE- QUENCY (%)	INDIVIDUAL WEIGHT (g)	TOTAL BIOMASS (%)
Aves			
Columbiformes			
Columbidae			
<i>Zenaida auriculata</i>	17.9	134.6	15.0
Passeriformes			
Emberizidae			
<i>Sicalis luteola</i>	1.8	16	1.8
<i>Zonotrichia capensis</i>	23.2	22.5	3.3
Mammalia			
Rodentia			
Caviidae			
<i>Cavia aperea</i> (adult)	3.6	630	7.0
<i>Cavia aperea</i> (young)	1.8	315	7.0
Muridae			
<i>Akodon azarae</i>	1.8	21	0.2
<i>Calomys musculus</i>	3.6	10	0.2
<i>Oryzomys flavescens</i>	1.8	17	0.2
<i>Rattus</i> spp.	43.0	250	66.9
Insecta			
Coleoptera			
Scarabaeidae			
<i>Sulcophanaeus menelas</i>	1.8	1	0.0

The Striped Owl is typically found in woodlands, forests, and savannas of tropical and subtropical zones (Grossman and Hamlet 1964). Our data show that it also occurs in the temperate-warm zone that corresponds to the Pampean Fitogeographic (Chaqueño Dominion, Cabrera 1976). In the past, this zone was dominated by tall grasslands without trees. Perhaps Striped Owls occur here because there is a natural corridor of tala forest which extends from the Entre Rios Province to Mar Chiquita Lagoon (Vervoorst 1967).

RESUMEN.—Se presenta información sobre la dieta del Lechuzón Orejudo (*Asio clamator*) en base al análisis de pellets (*N* = 34) y restos presa (*N* = 3), en el extremo Sur de su distribución Provincia de Buenos Aires, Argentina. Se identificaron 56 items presa, correspondiendo el 55.4% a los mamíferos, el 42.9% a las aves y el 1.8% a los insectos. El rango de pesos presa consumidos por esta lechuza fue de 1g a 630 g. El ítem mejor representado tanto en número (43%) como en biomasa (66.9%) fue *Rattus* spp. seguido en importancia numérica por el Chingolo Común (*Zonotrichia capensis*, 23.2%) y la Torcaza (*Zenaida auriculata*, 17.9%). A nivel de las aves el mayor aporte de biomasa fue dado por *Z. auriculata* (15%).

[Traducción de Autores]

ACKNOWLEDGMENTS

We thank R. Cáceres, M. Favero, S. and L. Bachmann, and students for field assistance, and A. Malizia for assisting in mammal identification. F. Jaksic and two anonymous referees made important contributions to improve this paper. Finally, we would like to express our especial gratitude to Mariano Manuel Martínez, and this paper is dedicated to his memory.

LITERATURE CITED

BLEDINGER, P., E. DE LUCA, AND M. SAGGESE. 1987. Nidificación otoño-invernal del Lechuzón Orejudo. *Nuestras Aves* 5:19.
BURTON, J. [ED.]. 1973. Owls of the world. Their evolution, structure and ecology. E.P. Dutton and Co., Inc., New York, NY U.S.A.
CABRERA, A. 1976. Regiones Fitogeográficas Argentinas. Enciclopedia Argentina de Agricultura y Jardinería. Tomo II, fascículo 1. Ed. Acme, Buenos Aires, Argentina.
CAMPERI, A.R. 1992. Estudio sobre aves colectadas en el extremo sudoeste de la Provincia de Buenos Aires. *Neotropica* 38:127–140.
CANEVARI, M., P. CANEVARI, G.R. CARRIZO, G. HARRIS, J. RODRIGUEZ MATA, AND R. STRANECK. 1991. Nueva guía de las aves argentinas. Tomos I y II. Fundación Acindar, Buenos Aires, Argentina.
GROSSMAN, M.L. AND J. HAMLET. 1964. Birds of prey of the world. Bonanza Books, New York, NY U.S.A.
HOLT, D.W., R. BERKLEY, C. DEPPE, P.L. ENRIQUEZ ROCHA,

P.D. OLSEN, J.L. PETERSEN, J.L. RANGEL SALAZAR, K P SEGARS, AND K.L. WOOD. 1999. Family Strigidae (Typical Owls). Pages 152–242 in J. del Hoyo, A. Elliot, and J. Sargatal [EDS.], Handbook of the birds of the world. Vol. 5. Barn Owls to hummingbirds. Lynx Edicions, Barcelona, Spain.
JIMÉNEZ, J.E. 1993. Notes on the diet of the Aplomado Falcon (*Falco femoralis*) in northcentral Chile. *J. Raptor Res.* 27:161–163.
MARTÍNEZ, M.M., J.P. ISACCH AND F. DONATTI. 1996. Aspectos de la distribución y biología reproductiva de *Asio clamator* en la Provincia de Buenos Aires, Argentina. *Ornitología Neotropical* 7:157–161.
NAROSKY, T. AND A. DI GIACOMO. 1993. Las aves de la provincia de Buenos Aires: distribución y estatus. Asoc. Ornitológica del Plata, Vázquez Mazzini Ed y L.O.L.A., Buenos Aires, Argentina.
PHELPS, JR., W.H. AND R. MEYER DE SCHAUENSEE. 1994. Una guía de las aves de Venezuela. Gráficas Armitano, Caracas, Venezuela.
REDFORD, K.H. AND J.F. EISENBERG. 1992. Mammals of the neotropics. The Southern Cone. Vol. 2. Univ. Chicago Press, Chicago, IL U.S.A.
SALVADOR, S.A. 1988. Datos de peso de aves argentinas. *Hornero* 13:78–83.
VERVOORST, F. 1967. Las comunidades vegetales de la depresión del Salado (Prov. de Buenos Aires). INTA. *La vegetación de la República Argentina* 7:1–262.

Received 1 May 1999; accepted 20 March 2000

J. Raptor Res. 34(3):237–241

© 2000 The Raptor Research Foundation, Inc.

DIET OF BREEDING CINEREOUS HARRIERS (*CIRCUS CINEREUS*) IN SOUTHEASTERN BUENOS AIRES PROVINCE, ARGENTINA

MARÍA S. BÓ, SANDRA M. CICCHINO, AND MARIANO M. MARTÍNEZ¹

Departamento de Biología, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata, Funes 3350, (7600) Mar del Plata, Argentina

KEY WORDS: *Cinereous Harrier*, *Circus cinereus*; *breeding diet*, *trophic niche breadth*; *Argentina*.

The Cinereous Harrier (*Circus cinereus*), one of two South American harriers, is widespread and distributed from Colombia and Ecuador, through Perú, Bolivia and Paraguay, southwestern Brazil to Tierra del Fuego and Islas Malvinas (Grossman and Hamlet 1964, Canevari et al. 1991, del Hoyo et al. 1994). In Argentina, it is most

common in Patagonia and Islas Malvinas (Narosky and Yzurieta 1987) but it has also been recorded throughout northwestern, central and, occasionally, the northeastern parts of the country (Canevari et al. 1991). The Cinereous Harrier inhabits savannas, grasslands, wetlands, marshes, lagoons, shrubsteppes, and shrublands 0–4500 m elevation (Jiménez and Jaksic 1988, Canevari et al. 1991, Narosky and Di Giacomo 1993, del Hoyo et al. 1994).

Little has been reported about the Cinereous Harrier. The few previous studies of this species have focused on aspects of ecology and behavior (Jiménez and Jaksic

¹ Deceased.